



Reay, M., Yates, C., Johnes, P., Arthur, C., Jones, D., & Evershed, R. (2019). High resolution HPLC-MS confirms overestimation of urea in soil by the diacetyl monoxime (DAM)colorimetric method. *Soil Biology and Biochemistry*, 135, 127-133.
<https://doi.org/10.1016/j.soilbio.2019.04.015>

Peer reviewed version

License (if available):
CC BY-NC-ND

Link to published version (if available):
[10.1016/j.soilbio.2019.04.015](https://doi.org/10.1016/j.soilbio.2019.04.015)

[Link to publication record in Explore Bristol Research](#)
PDF-document

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>




Fig. 1 Calibration curves for (a) allantoin, (b) alloxan, (c) hydantoin, and (d) biuret under the DAM colorimetric reaction conditions at concentrations between 0.1 to 10 mg N l⁻¹.




Fig. 2 Extracted ion chromatograms for urea [M+H]⁺ (m/z 61.08) for a soil extract from Merddwr, Conwy with (a) no added urea spike, and (b) 0.4 mg l⁻¹, (c) 2.0 mg l⁻¹ and (d) 4.0 mg l⁻¹ added urea spikes used for quantification by standard addition and (e) the standard addition calibration curve used to determine the urea concentration in the soil extract (0.709 $\mu\text{g g}^{-1}$).




Fig. 3 (a) Ureido group responsible for the cross-reactivity with the DAM colorimetric method, and environmentally relevant compounds containing the ureido group which cross-react: (b) allantoin; (c) hydantoin and (d) biuret.